

“ $\Omega > 1$ ”

“Sky-Notes” of the Open University Astronomy Club.

January 2019.

A Very Happy New Year to all readers!

Recent Events.

If you have any images and/or reports of recent events please contact Sheridan so that he can put them on the Club website.

If you wish to present them at a “Clubnight” meeting please contact Sheridan or myself before the meeting starts.

My apologies for the wrong date for the Mars – Neptune conjunction given in the December “Sky-Notes”. It should have read 7th not 14th. This was noticed by one of our members before the event and a suitable e-mail circulated.

Highlights of the Month.

1 st – 3 rd	Conjunctions between Moon, Venus and Jupiter in predawn SE sky.
2 nd	Saturn in Conjunction with Sun.
3 rd	Peak of Quadrantids meteor shower.
3 rd	Earth at Perihelion.
21 st	Total Eclipse of the Moon with the Moon at Perigee.
21 st – 23 rd	Conjunction between Venus and Jupiter in SE predawn sky.
30 th	Mercury at Superior Conjunction.
30 th – 31 st	Conjunctions between Moon, Venus and Jupiter in predawn SE sky.
Mars.	A moderately bright object in the S to W evening sky.
Jupiter .	Becoming better placed for “early hour” observation and imaging.
Saturn .	Reappearing low in SE dawn sky in the second half of the month.
Uranus.	Well placed for evening observation.
Neptune.	An early evening object low in SW.
Comet 36P/Wirtanen	Fading but very well placed for N observers.

Forthcoming Meetings.

OUAC.

Please note that the next “Clubnight” is on **Tuesday 8th January.**

BAA.

Full details of BAA meetings at: www.britastro.org

Other.

“European Astrofest 2019.”

Friday 8th and Saturday 9th February.

Kensington Conference & Events Centre, London.

A few minutes walk from High Street Kensington Underground Station.

Details at: europeanastrofest.com and the current issue of *Astronomy Now*.

Software.

A very useful item of Planetarium software is “Stellarium” and it’s FREE! Go to the website and download it and the associated user manual.

1. The Solar system.

Note all times shown are UT.

Earth.

Perihelion 03^d 05^h 20^m 147,099,766 km.

Aurora.

Long hours of darkness improve the opportunity for observing potential aurora.

Keep tuned to the www.spaceweather.com site for updates.

Subscribe (free) to the UK AuroraWatch website to receive alerts.

ISS.

Continues a series of morning passes during the first half of the month. Go to the “spaceweather” website and click the “Flybys” button and follow the instructions to set-up forecasts for your location. Alternatively go to the “Heavens Above” website and set-up for your location. Add to your “favourites”.

Sunrise and Sunset.

Bedford.

Latitude 52° 6.9’N Longitude 0° 28.1’W

Date.	Rise.	Transit.	Set.
01	08 ^h 11 ^m	12 ^h 05 ^m	16 ^h 00 ^m
08	08 ^h 09 ^m	12 ^h 09 ^m	16 ^h 09 ^m
15	08 ^h 04 ^m	12 ^h 11 ^m	16 ^h 19 ^m
22	07 ^h 57 ^m	12 ^h 14 ^m	16 ^h 31 ^m
29	07 ^h 47 ^m	12 ^h 15 ^m	16 ^h 43 ^m

Produced using Starry Night Pro v.6.

The Sun.

To prevent permanent damage to your eyes avoid looking at the Sun directly and never with binoculars or a telescope unless special (expensive!) filters are used.

The safest way is the simplest – project the image of the Sun onto grey or white card.

Take additional care if your telescope has plastic components. Plastic melts!

Very low activity at present (late Dec).

If you have or have access to observe in h-alpha the rewards are much greater.


Keep in touch with the Solar Dynamics Observatory satellite at <http://sdo.gsfc.nasa.gov/>

Add the “Spaceweather” and the “Soho Lasco C3” websites to your “favourite” websites.

The Moon.

Phases:

New	06 ^d 01 ^h 29 ^m	
First Quarter	14 ^d 06 ^h 47 ^m	
Full	21 ^d 05 ^h 17 ^m	Total Lunar Eclipse at Perigee. A ”Super Blood Moon”.
Last Quarter	27 ^d 11 ^h 29 ^m	



The Moon cont.

Apsides:

Apogee	09 ^d 04 ^h	Diameter. 29' 25"	Distance. 406,113km.
Perigee	21 ^d 20 ^h	Diameter. 33' 26"	Distance. 357,344km.

For northern observers:

The waxing crescent Moon is becoming better placed.

The waxing gibbous Moon is very well placed.

The Full Moon is very well placed.

The waning gibbous Moon is well placed.

The waning crescent Moon is less well placed.

Observe the regions along the terminator (sunrise and sunset on the Moon) where the low angle of the Sun highlights lunar topography. A basic lunar map is all you need to get started. *Sky & Telescopes* "Lunar 100 Card" is another good starting point. If you are starting out on photography and/or imaging the Moon provides an excellent target.

Lunar Occultations.

Unlike the gradual disappearance of a planet (small disc) a star vanishes instantly demonstrating that it is a point source of light as viewed from the earth. For all occultation events start observing 10 to 15 minutes before the predicted time to identify the required star and to allow for slightly different time if you are not at Greenwich. Use an accurate watch to record the time that *you* observe the occultation remembering that times are UT not BST. Enter details in your observing log.

Further details of occultations can be found in current *BAA Handbook* and monthly periodicals such as *Astronomy Now* and *Sky at Night*.

Observing Opportunities and Challenges!

On 4th and 5th try locating the very thin crescent Moon very low in the ESE dawn skies **before sunrise**.

On 7th and 8th try locating the very thin crescent Moon in SW evening twilight **after sunset**.

If you can take images of the above so much the better.

The Planets.

The predawn and dawn SE sky provides the greatest interest this month with pleasant conjunctions between the Moon and planets at the beginning and end of the month.

Mercury.

Difficult to spot very low in the SE dawn twilight during first half of the month.

Superior Conjunction 30th.

Moon close on 4th.

Date.	Mag.	Dia.	Phase.	Rise.	Transit.	Set.
01	-0.4	5.2''	0.90	07 ^h 04 ^m	10 ^h 55 ^m	14 ^h 46 ^m
15	-0.7	4.8''	0.97	07 ^h 47 ^m	11 ^h 33 ^m	15 ^h 20 ^m

Venus.

Dominates the SE predawn/dawn sky.

Greatest Elongation W (47°) on 6th.

Conjunction with Jupiter on 21st – 23rd.

Pleasant conjunctions with Jupiter and the Moon on 30th and 31st.

Moon close on 1st, 2nd and 31st.

Date.	Mag.	Dia.	Phase.	Rise.	Transit.	Set.
01	-4.5	26''	0.48	04 ^h 07 ^m	08 ^h 48 ^m	13 ^h 29 ^m
31	-4.3	19''	0.62	04 ^h 55 ^m	09 ^h 03 ^m	13 ^h 11 ^m

Mars.

The disc is rapidly shrinking making surface details more difficult to observe and/or image.

However increasing N declination improves seeing conditions.

Moon close on 12th.

Date.	Mag.	Dia.	Phase.	Rise.	Transit.	Set.
01	+0.5	7.4''	0.87	11 ^h 17 ^m	17 ^h 20 ^m	23 ^h 23 ^m
31	+0.9	6.2''	0.89	09 ^h 49 ^m	16 ^h 36 ^m	23 ^h 23 ^m

The Mars **Curiosity** rover continues to return excellent data and images.

Contact has yet to be re-established with the **Opportunity** rover.

Mission details and progress are on the appropriate NASA websites.

Jupiter.

Becoming better placed for observation and/or imaging low in the SE predawn sky.

Conjunction with Venus on 21st – 23rd.

Pleasant conjunction with Venus and the Moon on 30th and 31st.

Moon close on 11th.

Excellent target for imaging.

See BAA *Handbook* and/or monthly periodicals for satellite phenomena.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
01	-1.8	32''	05 ^h 59 ^m	10 ^h 01 ^m	14 ^h 02 ^m
31	-1.9	34''	04 ^h 27 ^m	08 ^h 27 ^m	12 ^h 25 ^m

Saturn.

Reappearing low in the SE predawn sky towards end of the month.

Rings wide open.

Moon close on 5th.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
15	+0.5	15''	07 ^h 23 ^m	11 ^h 20 ^m	15 ^h 17 ^m
31	+0.6	15''	06 ^h 27 ^m	10 ^h 25 ^m	14 ^h 48 ^m

Uranus.

Well placed for evening observation.

Moon close 14th.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
01	+5.8	3.6''	12 ^h 07 ^m	19 ^h 04 ^m	02 ^h 06 ^m
31	+5.8	3.5''	10 ^h 09 ^m	17 ^h 07 ^m	00 ^h 10 ^m

Neptune.

Sinking into the SW early evening skies.

Moon close 10th.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
01	+7.9	2.2''	10 ^h 55 ^m	16 ^h 21 ^m	21 ^h 46 ^m
31	+8.0	2.2''	08 ^h 58 ^m	14 ^h 26 ^m	19 ^h 53 ^m

Dwarf Planets.

Ceres.

An 8th mag object located in Libra. About 5° NW of Venus at the start of the month. Forms a pleasant triangle with the crescent Moon to the west on 1st.

Eris.

A CCD target object in Cetus.

Haumea.

A CCD target located in Bootes.

MakeMake.

A CCD target in Coma Berenices.

Pluto.

In conjunction with the Sun on 11th. Not observable.

Asteroids. (Approx mag +10.5 or brighter).

- Vesta (4).** A 7th mag object located in Capricornus. Becoming lost in the SW evening twilight.
- Pallas (2).** Located in Virgo. Brightening from +9.0 to +8.6.
- Juno (3)** Moving from Eridanus into Taurus. Mag +8.2 at opposition on 2nd.
- Hebe (6).** Located in Orion. Mag +8.5 fading to +9.2
- Eros (433).** Moving from Perseus into Taurus. Mag +9.1 fading to +9.3.
- Herculina (532).** Located in Leo. Brightening from +9.6 to +8.9.

Charts and details of asteroids one month either side of opposition are available at:

http://britastro.org/computing/charts_asteroid.html

See also the *BAA Handbook* and/or monthly periodicals.

Comets.

Comet 46P/Wirtanen.

Fading from 5th to 7th magnitude as it tracks through Lynx into Ursa Major.

Comet 38P/Stephan -Oterma.

Fading from 10th to 11th magnitude as it tracks through Lynx.

Comet 64P/Swift-Gehrels.

An 11th magnitude object tracking from Aries into Taurus. It will be track about 4° N of the Pleiades between 18th to 25th.

Useful articles for the above in the January issues of *Astronomy Now* and *Sky at Night.s*

Charts and details of selected comets are available at:

http://britastro.org/computing/charts_comet.html

See also the *BAA Handbook* and/or monthly periodicals.

Meteor Showers.

The **Quadrantids** are active from 1st – 6th January with peak activity on 4^d 03^h, ZHR = 80+. Very favourable as no interference from moonlight!

There are always **sporadic** events and the chance of a brilliant fireball. The latter should be recorded and reported.

Near Earth Objects.

Please refer to www.spaceweather.com for updates.

Eclipses.

A **Total Lunar Eclipse** occurs on 21st visible from UK. A “Super Blood Moon”!

Penumbral eclipse commence (P1) at 02:37.

Umbral eclipse commences (U1) at 0:34.

Total eclipse commences (U2) at 04:41.

Greatest eclipse occurs at 05:12.

Total eclipse ends (U3) at 05:43.

Umbral eclipse ends (U4) at 06:51.

Penumbral eclipse ends (P4) at 07:48.

Everything crossed for a clear sky!

2. The Deep Sky.

Abbreviations used.

M = Messier object (Shown in **bold**).

NGC = New General Catalogue. IC = Index Catalogue (Extension of the NGC).

ds = double star. ts = triple star. ms = multiple star. vs = variable star.

gc = globular cluster. oc = open cluster. pn = planetary nebula.

en = emission nebula. rn = reflection nebula. sg = spiral galaxy.

eg = elliptical galaxy. lg = lenticular galaxy. ir = irregular galaxy.

pg = peculiar galaxy. snr = super nova remnant. ly = light year.

The magnitude of an object, excluding double, triple, multiple and variable stars, is shown in brackets e.g. (6.5).

All magnitudes are + unless otherwise shown.

2.1 Variable Stars of the month.

Beta (β) Persei, Algol. Range 2.2 to 3.4, period 2.7 days. Becoming well placed by mid evening. Favourable minima at “social hours” occur on 18^d 0.00^h, 20^d 20.8^h and 23^d 17.6^h.

Delta (δ) Cephei. Range 3.5 to 4.4, period 5.37 days. The prototype for the Cepheid class of variable stars. Their period-luminosity relationship has led them to being used as “standard candles” in measuring distances to nearby galaxies.

Mu (μ) Cephei. Range 3.7 to 5.0, approximate period 755 days. A semi-regular variable star famous for its striking red colour being fittingly called “Herschel’s Garnet Star”. It is the reddest naked eye star visible from the northern hemisphere. Its colour may show signs of variability.

Omicron (\omicron) Ceti Mira. The classic long period variable star fading from maximum (+3.4) towards minimum in Jun/Jul.

U Ori. Well placed for nightlong observation. Brightening towards (+6.3) in Apr/May 2019.

2.2 Double Stars of the month.

Beta Cam. See notes below.

Alpha Gem (Castor). See notes below.

Delta Gem. See notes below.

Gamma Lep. See notes below.

Beta Mon. See notes below.

Beta Ori (Rigel). See notes below.

Sigma Ori. See notes below.

Theta-1 Ori (The Trapezeium). See notes below.

Struve 742 Tau. See notes below.

2.3 This Month's Constellations - Double Stars/Star Clusters/Nebulae/Galaxies.

Auriga (Aur).

NGC1960 (M36) (6.0) oc. Large bright grouping. In same low power field as M38.

NGC2099 (M37) (5.6) oc. Richest and brightest of the three Messier star clusters in Auriga. At 150 stars brighter than 12th magnitude.

NGC1912 (M38) (6.4) oc. Larger than M36. Many bright stars arranged in pairs. The above are excellent objects for photography. Guided exposures of a few minutes will be necessary. CCD images require much shorter exposures.

NGC1664 (7.6) oc. Fine cluster on the border of Auriga and Perseus.

NGC1778 (7.7) oc. A 6" telescope will show about 20 stars. Larger apertures will reveal more.

NGC1857 (7.0) oc. Hazy patch surrounding an orange 7th magnitude star which interferes with viewing the fainter stars.

NGC1893 (7.5) oc. Fine, though rather sparse cluster. 8"+ telescopes under dark skies may begin to reveal the pale light of the brightest part of the emission nebula IC410 which pervades the star cluster.

NGC1907 (8.2) oc. This small cluster lies just west of M38 appearing as a small smudge of light.

NGC1931 (11.3) en. An 8" telescopes from dark skies should reveal this small pea-nut shaped emission nebula.

NGC 2192 (10.9) oc. Not an easy object probably requiring a 6" telescope to locate and 10"+ to resolve.

NGC2281 (5.4) oc. Handful of stars often overlooked.

IC405 en (6v) The "Flaming Star Nebula". Illuminated by the star AE Aur which is a "runaway star" whose path can be traced back to Orion. At present the star is passing by/through the normally dark dust and gas cloud of IC405 and thus illuminating it. In the future as AE moves away the nebula will again become dark.

Camelopardalis (Cam).

Beta (β) ds. (4.0/8.6, sep. 80.8"). Wide yellow and blue pair (AB). From dark sites the system can be seen to be embedded in a streak of dark nebulosity. A third magnitude 11.8", sep (BC) 14.8", component is visible in moderate apertures.

1 ds. (5.7/6.8, sep. 10.3"). White and pale blue pair.

11 & 12 ds. (5.4/6.5, sep, 108.5") White pair. Easy object for binoculars and small telescopes.

29 ds. 6.5/9.5, sep. 25.1". Yellow primary with fainter pale blue secondary.

Struve (Σ)390 ds. 5.1/9.5, sep. 14.8". White primary with fainter purple secondary.

Struve (Σ)1122 ds. (7.8/7.8, sep. 15.4"). Fine pair of equally bright white stars.

Struve (Σ)1625 ds. 7.3/7.8, sep. 14.4". Almost equal pair of yellowish stars.

Struve (Σ)1694 ds. 5.3/5.8, sep.21.6". Easy pair of whitish stars.

NGC1501 (11.5) pn. Bright, large with a blue tint located about 2 degrees south of the open star cluster NGC1502. Resembles the "Eskimo Nebula" in Gemini. NGC1502 (5.7) oc. A fine open cluster. Extending NW of this cluster is a line of 9th and 10th magnitude stars that form "Kemble's Cascade". A beautiful sight in binoculars or a low power wide field eyepiece on small telescopes.

NGC2403 (8.4) sg. Visible in large binoculars. It lies at a distance of 8m light years and is possibly a member of the M81/M82 group.

IC342 (12.0) sg. Once considered to be a member of the Local Group of galaxies. Its low surface brightness makes it a challenge for moderate apertures. This face on spiral galaxy would be an impressive object if it were further away from the plane of the Milky Way.

Gemini (Gem).

Alpha (α) **Castor** ms. 1.9/2.9 sep 4.0". Close visual pair. However each of these is a spectroscopic binary. A more distant ninth magnitude star (red) forms part of an eclipsing binary system. A fascinating family!

Delta (δ) ds. 3.5/8.2 sep 5.8". Yellow primary with bluish secondary.

Kappa (κ) ds. 3.6/8.1 sep 7.1". Orange-yellow primary with bluish companion.

Lambda (λ) ds. 3.6/10.7 sep 9.6". Blue-white primary with bluish companion.

Σ 1108 (Struve) ds. 6.6/8.3 sep 11..5". Yellow primary with bluish companion.

M35 (5.1) oc. Just visible to the naked eye from dark sites. It is a superb object in telescopes. On its western edge lies the more distant open star cluster IC2158.

NGC2129 (10.2) oc. Located about a degree SW of IC2158.

NGC2266 (9.5) oc. Located about two degrees north of ϵ Gem.

NGC2392 (10.5) pn. The "Eskimo nebula" is a fine planetary nebula located about two degrees SE of δ . The nickname is derived from the appearance of a face surrounded by the hood of a parka.

NGC2420 (8.3) oc. Located about two degrees east of the "Eskimo".

Complete this deep-sky tour of Gemini by locating the open star clusters NGC2355 (9.7) and NGC2395 (7.1).

Lepus (Lep).

Lying beneath Orion Lepus is easily recognized by a quadrilateral of four third magnitude stars and contains a variety of deep-sky objects including one Messier object.

Alpha (α) ds; (2.6,11.1; sep. 35.8").

Beta (β) ds; (3.0/7.5; sep. 2.3").

Gamma (γ) ds. (3.7/6.3, sep. 96.3"). Fine yellow and pale orange pair.

Kappa (κ) ds; (4.5/7.4; sep. 2.6"). White and blue companions.

Iota (ι) ds; (4.5/10.8; sep. 12.7").

NGC1974 sg (11.8). Seen almost edge on.

NGC1904 (**M79**) gc (8.0). A fine globular cluster visible as a fuzzy spot in binoculars. Outer edges begin to resolve in 12" (30cm) telescopes.

IC418 pn (10.7). Very small but bright. Central 10.7 mag star surrounded by pale ring. Use a UHC or OIII filter for best results.

Monoceros (Mon).

This faint and rather indistinct constellation is located between Orion and Canis Minor.

Beta (β) ts. 4.7/5.2/6.1 Sep. AB = 7.3', sep BC = 2.8". Striking triple of bluish white stars.

Epsilon (ϵ) ts. 4.5/6.5 sep. 13.4". Close pair of pale yellow stars. The third mag 12.7 bluish white member is visible in 12"+ apertures.

NGC2244 oc (4.8). Fine open star cluster surrounded by NGC2237-9 "The Rosette Nebula" which is best seen using a UHC filter. Shows well in photographs.

NGC2261 en (10v). "Hubble's Variable Nebula". Located about 2^o southwest of NGC2264 this a fascinating object and well worth monitoring for changes in shape and brightness due to the enveloped variable star R Monocerotis. The triangular wedge appears is almost comet like. Detailed star chart available for telescope owners.

NGC2264 oc + en (4.0) The "Christmas Tree Cluster". A fine open cluster with associated nebula that includes the "Cone Nebula".

NGC2323 (**M50**) oc (5.9). Superb open cluster.

There are many other open clusters in this area of the Milky Way - NGC's 2215(8.4), 2286(7.5), 2301(6.0), 2335(7.2), 2343(6.7), 2353(7.1) and 2506(7.6).

Orion (Ori).

This constellation dominates the winter skies and because it is so easily recognized forms one of the "key constellations" for finding other winter groupings.

Orion's two brightest stars provide a marked contrast. Betelgeuse is distinctly orange in colour. It is a red giant star entering old age. Rigel is a brilliant blue/white star indicating the exuberance of youth. Betelgeuse is slightly variable in brightness, range 0.1 - 0.9 and bears the designation α (alpha) indicating that it was brighter than Rigel, β (beta) (0.1), when stars were given these designations. Rigel is now the brighter of the two so either early magnitude estimates were wrong or Betelgeuse has dimmed slightly.

Beta (β) Rigel ds. 0.1/6.8 sep. 9.5". Brilliant bluish white primary with much fainter bluish secondary.

Eta (η) ds. 3.6/5.0 sep. 1.5". Close pair of white stars.

Delta (δ) ds. 2.0/6.9 sep. 52.6". Blue white primary with pale blue secondary.

Lambda (λ) ds. 3.5/5.6 sep.4.4". White stars. Part of a quadruple system.

Theta-1 (θ) ms. "The Trapezium". AB: 6.7/7.9 sep. 8.8"; CD: 5.1/6.7 sep. 13.4". Superb object!

Iota (ι) ts. 2.8/7.3 sep. 11.3". White primary with pale blue secondary. The third reddish 11th magnitude component is located 50" away.

Sigma (σ) ms. 4.0/10.3 sep. 11.4. Colourful multiple star. Fainter triple star Σ 761 in same field.

Zeta (ζ) Alnitak ds. 1.9/4.0 sep. 2.3". Bluish white stars. Part of a triple system with the Flame Nebula (NGC2024) in the field of view.

NGC1976 (**M42**) en. One of the most famous objects in the sky. Marking Orion's sword the "Great Orion Nebula" is visible to the naked eye as a faint misty patch. A pair of binoculars or small telescope will begin to reveal detail. Increasing aperture and low power bring increasing rewards for the visual observer. Embedded in the nebula is **Theta (θ) Ori**. A group of four young stars, mags 5.4,6.3, 6.8 and 7.0, aptly called "The Trapezium". The whole nebula is a stellar nursery with spectacular images being obtained from large Earth based telescopes and the Hubble Space Telescope. M42 is an ideal target for photography.

NGC1982 (**M43**) en. A small patch of nebulosity on the northern edge of M42.

NGC2024 (en), nicknamed "the Flame Nebula", surrounds ζ Ori.

IC434 en is a strip of nebulosity just south of ζ . The famous "Horse's Head Nebula" (Barnard 33) is a small dark intrusion seen dramatically in photographs. It provides one of the biggest challenges to visual observers requiring very dark transparent skies. Responds well to a H-beta nebula filter.

NGC2068 (**M78**) (8.0) rn is a small patch of nebulosity about two degrees NNE of ζ .

NGC2112 (9.1) oc is an open star cluster about two degrees east of M78.

Other open clusters worth locating are NGC2186 (8.7), NGC2169 (5.9) and NGC2175 (6.8) which superimposes a small patch of nebulosity NGC2174.

Long exposure photographs reveal a long arc of nebulosity curving up the east side Orion. This is called "Barnard's Loop" which is extremely difficult to discern visually almost regardless of aperture. Remarkably it has been seen with the naked eye (initially by accident!) from dark sites using O III or UHC filters. The "Loop" is a faint ring of hot gas some 14^o by 10^o with the western part of the ring being less distinct. The "ring" may be due to radiation pressure from the hot young stars in the region of Orion's belt/sword acting on interstellar material. A less favoured school of thought is that it may be a supernova remnant.

Taurus (Tau).

Lambda (λ) vs. Eclipsing binary, range 3.4 to 4.1, period 3.95d.

Phi (ψ) ds. 5.0/8.4 sep 52.1". Fine contrasting deep yellow primary with blue companion.

Σ 495 (Struve) ds. 6.0/8.8 sep 3.8". Fine pair of yellow stars.

Σ 742 ds. 7.1/7.5 sep 4". Fine pair in the same wide field as the Crab Nebula (M1) to the E. Often overlooked when locating M1.

47 Tauri ds. 4.9/7.4 sep 1.1". Very close pair of yellow stars.

T vs. Irregular variable, range 9.4 to 13.5, period erratic.

RV vs. Irregular variable, range 9.5 to 13, period 79d.

RR vs. Irregular variable, range 9.9 to 13, period "chaotic".

M45 - "The Pleiades" or "Seven Sisters" oc. Probably the most famous star cluster. Test your eyesight from a dark site by counting the number of naked eye stars that are visible. Seven should readily be seen. Keen vision will lead you into double figures. A test for moderate apertures is the nebulosity around some of the other brighter stars of the group, especially Merope. Nebula/UHC filters will help.

The Hyades oc. Another naked eye star cluster although more spread out than the Pleiades. The first magnitude star Aldebaran (0.9) is not a member of the cluster and appears brighter as it is closer to us, 21 parsecs - about halfway to the clusters centre. Aldebaran has a faint companion (13.4) separation 30.4". Spotting the companion is not easy even in moderate telescopes.

NGC1514 (10.0) pn. Not an easy object in small instruments. An OIII filter will help.

NGC1554/1555 is a faint variable reflection nebula illuminated by the very young star T Tauri (see above).

NGC1647 (6.4) oc. Rich cluster.

NGC1746 (6.1) oc. Poorer than 1647

NGC1808/1817 (7.0/7.7) oc. a pleasant "double cluster" when observed at low powers.

NGC1952 (**M1**)(8.5) snr. "The Crab Nebula". One of the most studied astronomical objects in recent decades. A "new star" appeared in 1054 and over a few months faded from view. Centuries later the faint oval patch was discovered by Dr John Bevis in 1731 and independently by Charles Messier on 12th September 1758 while observing the comet of that year. Its true nature was not recognized until much later. In 1942 astronomers had speculated that a peculiar star in the nebula might be a neutron star. In 1968 radio astronomers discovered that the central star was a pulsar spinning at 30 times a second i.e. the period of the pulses is 33 milliseconds. The "Crab" is also a strong emitter of X-rays.

Happy and Successful Observing!

P.V.H.