

“ $\Omega > 1$ ”

“Sky-Notes” of the Open University Astronomy Club

January 2025

**A Very Happy and Successful New Year and
Best Wishes for 2025 to all readers of “ $\Omega > 1$ ”!**

Recent Events

If you have any images and/or reports of recent events please contact Sheridan so that he can put them on the Club website

If you wish to present them at a “Clubnight” meeting please contact Adrian or myself before the meeting starts

Highlights of the Month

3 rd	Peak of Quadrantids meteor shower
4 th	Earth at Perihelion
4 th	Lunar Occultation of Saturn See notes below
10 th	Venus at Greatest Elongation East
12 th	Mars closest to the Earth
16 th	Mars at Opposition
25 th	Burns’ Night
Mercury	Low in the SE morning twilight at the beginning of the month
Venus	Brilliant object in the SW evening sky
Mars	See notes below
Jupiter	Well placed for evening through midnight observation
Saturn	Early evening object in the SW sky
Uranus	Well placed for evening through midnight observation
Neptune	Sinking into the WSW evening sky
Asteroids at Opposition – see notes below	
Comets and Meteor Showers – see notes below	

Forthcoming Meetings

OUAC

The next “Clubnights” will be held on **Tuesday 7th January** and **Tuesday 4th February**

Software

A very useful item of Planetarium software is “**Stellarium**” and it’s **FREE!**
Go to the website and download it and the associated user manual

1 The Solar system

Note all times shown are UT

Earth

Perihelion 04^d 13^h 28^m 147,103,682 km

Aurora

Long hours of darkness improve the opportunity for observing potential aurora.

Keep tuned to the www.spaceweather.com site for updates.

Subscribe (free) to the UK AuroraWatch website to receive alerts.

Artificial Satellites.

For the ISS and other “bright” satellites go to the “Heavens Above” website and set-up for your location.

Alternatively go to “spaceweather” website and click the “Flybys” button and follow the instructions to set-up forecasts for your location.

Add to your “favourites”.

Sunrise and Sunset

Bedford

Latitude 52° 6.9’N Longitude 0° 28.1’W

Date.	Rise.	Transit.	Set.
01	08 ^h 11 ^m	12 ^h 06 ^m	16 ^h 01 ^m
08	08 ^h 08 ^m	12 ^h 09 ^m	16 ^h 09 ^m
15	08 ^h 03 ^m	12 ^h 12 ^m	16 ^h 20 ^m
22	07 ^h 56 ^m	12 ^h 14 ^m	16 ^h 31 ^m
29	07 ^h 48 ^m	12 ^h 15 ^m	16 ^h 44 ^m

Produced using Starry Night Pro v6

The Sun.

To prevent permanent damage to your eyes avoid looking at the Sun directly and never with binoculars or a telescope unless special (expensive!) filters are used.

The safest way is the simplest – project the image of the Sun onto grey or white card.

Take additional care if your telescope has plastic components. Plastic melts!

If you have or have access to observe in h-alpha the rewards are much greater.

Solar Cycle 25 has reached **SolarMax** and should continue to provide good activity for the next couple of years.

Keep in touch with the Solar Dynamics Observatory satellite at <http://sdo.gsfc.nasa.gov/>

Add the “spaceweather” and “Soho” websites to your “favourite” websites.

The Moon

Phases:



Produced using Lunar Phase Pro

First Quarter	06 ^d 23 ^h 56 ^m
Full	13 ^d 22 ^h 26 ^m
Last Quarter	21 ^d 20 ^h 31 ^m
New	29 ^d 12 ^h 37 ^m

Apsides:

Perigee	08 ^d 00 ^h	Diameter. 32' 16"	Distance. 370,173km.
Apogee	21 ^d 05 ^h	Diameter. 29' 32"	Distance. 404,298km.

Observing/Imaging

For northern observers:

- The waxing crescent Moon is becoming better placed.
- The waxing gibbous Moon is very well placed.
- The Full Moon is very well placed.
- The waning gibbous Moon is well placed.
- The waning crescent Moon is less well placed.

The Moon cont

Observe the regions along the terminator (sunrise and sunset on the Moon) where the low angle of the Sun highlights lunar topography.

Take advantage of the prevailing Lunar Libration to observe features on the Limb that are normally hidden or are better presented.

A basic lunar map is all you need to get started. *Sky & Telescopes* "Lunar 100 Card" is another good starting point.

If you are starting out on imaging the Moon provides an excellent target.

On 1st (difficult!) and 2nd try locating the very thin crescent Moon in SW evening twilight **after sunset**.

On 27th and 28th (difficult!) try locating the very thin crescent Moon very low in the ESE dawn sky **before sunrise**.

Lunar Occultations

Unlike the gradual disappearance of a planet (small disc) a star vanishes instantly demonstrating that it is a point source of light as viewed from the earth. For all occultation events start observing 10 to 15 minutes before the predicted time to identify the required star and to allow for slightly different time if you are not at Greenwich. Use an accurate watch to record the time that *you* observe the occultation remembering that times are UT not BST. Enter details in your observing log.

Occultation of Saturn on 4th.

Timings are for Greenwich. Check exact times for your location.

Disappears behind the Dark Limb (DD)	17 ^h 21.3 ^m
Reappears from behind the Bright Limb (RB)	18 ^h 30.5 ^m

Details of occultations can be found in current *BAA Handbook* and monthly periodicals such as *Astronomy Now* and *Sky at Night*.

Simulate events for your location on "Stellarium" or similar software.

The Planets

Mercury

Bright object in the SE morning twilight in the first few days of the month but then becomes increasingly more difficult to spot during the month.

Do not sweep the area with binoculars or a telescope if sunrise is imminent!

Moon close on ?th.

Date.	Mag.	Dia.	Phase.	Rise.	Transit.	Set.
01	-0.4	5.8''	0.78	06 ^h 36 ^m	10 ^h 35 ^m	14 ^h 34 ^m
15	-0.4	5.0''	0.91	07 ^h 19 ^m	11 ^h 06 ^m	14 ^h 54 ^m

Venus

Brilliant object dominating the SW early evening sky.

Greatest Elongation East (47.2°) on 10th.

Dichotomy (50% phase) on 12th.

Then becomes a waning crescent with increasing apparent diameter.

2° north of Saturn on 20th.

Moon close on 3rd.

Date.	Mag.	Dia.	Phase.	Rise.	Transit.	Set.
01	-4.4	22''	0.55	10 ^h 27 ^m	15 ^h 20 ^m	20 ^h 14 ^m
10	-4.4	25''	0.51	10^h 04^m	15^h 20^m	20^h 35^m
31	-4.6	32''	0.38	08 ^h 58 ^m	15 ^h 04 ^m	21 ^h 11 ^m

Mars

Closest to the Earth on 12th.

Opposition on the 16th.

The “small” diameter at this Opposition is compensated by its high northern declination which benefits northern observers.

Moon close on 14th.

Date.	Mag.	Dia.	Phase.	Rise.	Transit.	Set.
01	-1.2	14''	0.99	17 ^h 13 ^m	01 ^h 38 ^m	09 ^h 58 ^m
16	-1.4	15''	1.00	15^h 38^m	00^h 16^m	08^h 47^m
31	-1.1	14''	0.99	14 ^h 08 ^m	22 ^h 48 ^m	07 ^h 32 ^m

Jupiter

Prominent object very well placed for evening through midnight observation.

Moon close on 10th.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
01	-2.7	47''	13 ^h 56 ^m	22 ^h 02 ^m	06 ^h 13 ^m
31	-2.5	43''	11 ^h 50 ^m	19 ^h 56 ^m	04 ^h 06 ^m

See BAA *Handbook* and/or monthly periodicals for satellite phenomena.

Simulate events on “Stellarium” or similar software.

Saturn

Sinking into the SW early evening sky.

2° S of Venus on 20th.

Titan shadow transit on 7th and 23rd. Simulate events on “Stellarium”.

Lunar Occultation on 4th. See notes above.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
01	+1.1	17”	11 ^h 00 ^m	16 ^h 22 ^m	21 ^h 44 ^m
31	+1.1	16”	09 ^h 06 ^m	14 ^h 34 ^m	20 ^h 02 ^m

Uranus

Located on Aries/Taurus border and well placed for evening through midnight observation.

Moon close on 9th.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
01	+5.7	3.7”	12 ^h 56 ^m	20 ^h 40 ^m	04 ^h 29 ^m
31	+5.7	3.6”	10 ^h 57 ^m	18 ^h 41 ^m	02 ^h 29 ^m

Neptune

Located in Pisces now sinking into the WSW evening sky.

Moon close 5th.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
01	+7.9	2.3”	11 ^h 16 ^m	17 ^h 08 ^m	22 ^h 59 ^m
31	+7.9	2.2”	09 ^h 19 ^m	15 ^h 12 ^m	21 ^h 05 ^m

Dwarf Planets

Ceres

Too close to the Sun for observation.

Eris

A CCD target object in Cetus.

Haumea

A CCD target located in Boötes.

MakeMake.

A CCD target in Coma Berenices.

Pluto

Not observable. **Conjunction with the Sun on 21st.**

Asteroids (Approx mag +10.5 or brighter).

Vesta (4) A 7th mag morning object on Virgo.

Irene (14) Located in Gemini. Mag +9.5 at **Opposition on 3rd.**

Alinda (887) Located in Gemini. Mag +9.2 at **Opposition on 12th.** Included as making a “close” approach to the Earth.

Charts and details of asteroids one month either side of opposition are available at:

http://britastro.org/computing/charts_asteroid.html

See also the “Heavens above” website, the *BAA Handbook* and/or monthly periodicals.

Comets

No “bright” comets this month.

The “Heavens above” website gives very useful details of comets currently available for observation/imaging.

Charts and details of selected comets are available at:

http://britastro.org/computing/charts_comet.html

See also the *BAA Handbook* and/or monthly periodicals.

Meteor Showers

The **Quadrantids** are active from 1st – 6th January with peak activity (in daylight) on 3rd 11^h - 14^h, ZHR = 80+. Rather narrow peak as indicated. Little interference from Moonlight either side of the peak.

The shower produces “bright events” so still worth observing.

There are always **Sporadic** events and the chance of a brilliant fireball. The latter should be recorded and reported.

Near Earth Objects

Please refer to www.spaceweather.com for updates.

Eclipses

No Lunar or Solar Eclipses this month.

2. The Deep Sky

Abbreviations used

M = Messier object (Shown in **bold**).

NGC = New General Catalogue. IC = Index Catalogue (Extension of the NGC).

ds = double star. ts = triple star. ms = multiple star. vs = variable star.

gc = globular cluster. oc = open cluster. pn = planetary nebula.

en = emission nebula. rn = reflection nebula. sg = spiral galaxy.

eg = elliptical galaxy. lg = lenticular galaxy. ir = irregular galaxy.

pg = peculiar galaxy. snr = super nova remnant. ly = light year.

The magnitude of an object, excluding double, triple, multiple and variable stars, is shown in brackets e.g. (6.5).

All magnitudes are + unless otherwise shown.

2.1 Variable Stars of the month

Beta (β) Persei, Algol.

Range +2.2 to +3.4, period 2.7 days. Becoming well placed by mid evening.

Favourable minima at “social hours” occur on 8^d 0.6^h, 10^d 21.5^h, 13^d 18.3^h and 30^d 23.2^h.

Delta (δ) Cephei.

Range 3.5 to 4.4, period 5.37 days. The prototype for the Cepheid class of variable stars.

Their period-luminosity relationship has led them to being used as “standard candles” in measuring distances to nearby galaxies.

Mu (μ) Cephei.

Range 3.7 to 5.0, approximate period 755 days. A semi-regular variable star famous for its striking red colour being fittingly called “Herschel’s Garnet Star”. It is the reddest naked eye star visible from the northern hemisphere. Its colour may show signs of variability.

U Ori.

Well placed for nightlong observation. Brightening from minimum (+12) in December 2024 towards maximum (+6.3) in June 2025.

2.2 Double Stars of the month

Beta Cam. See notes below.

Alpha Gem (Castor). See notes below.

Delta Gem. See notes below.

Gamma Lep. See notes below.

Beta Mon. See notes below.

Beta Ori (Rigel). See notes below.

Sigma Ori. See notes below.

Theta-1 Ori (The Trapezeium). See notes below.

Struve 742 Tau. See notes below.

2.3 This Month's Constellations - Double Stars/Star Clusters/Nebulae/Galaxies

Auriga (Aur)

NGC1960 (M36) (6.0) oc. Large bright grouping. In same low power field as M38.

NGC2099 (M37) (5.6) oc. Richest and brightest of the three Messier star clusters in Auriga with 150 stars brighter than 12th magnitude.

NGC1912 (M38) (6.4) oc. Larger than M36. Many bright stars arranged in pairs.

The above are excellent objects for photography. Guided exposures of a few minutes will be necessary. CCD images require much shorter exposures.

NGC1664 (7.6) oc. Fine cluster on the border of Auriga and Perseus.

NGC1778 (7.7) oc. A 6" telescope will show about 20 stars. Larger apertures will reveal more.

NGC1857 (7.0) oc. Hazy patch surrounding an orange 7th magnitude star which interferes with viewing the fainter stars.

NGC1893 (7.5) oc. Fine, though rather sparse cluster. 8"+ telescopes under dark skies may begin to reveal the pale light of the brightest part of the emission nebula IC410 which pervades the star cluster.

NGC1907 (8.2) oc. This small cluster lies just west of M38 appearing as a small smudge of light.

NGC1931 (11.3) en. An 8" telescopes from dark skies should reveal this small pea-nut shaped emission nebula.

NGC 2192 (10.9) oc. Not an easy object probably requiring a 6" telescope to locate and 10"+ to resolve.

NGC2281 (5.4) oc. Handful of stars often overlooked.

IC405 en (6v) **"The Flaming Star Nebula"**. Illuminated by the star AE Aur which is a "runaway star" whose path can be traced back to Orion. At present the star is passing by/through the normally dark dust and gas cloud of IC405 and thus illuminating it. In the future as AE moves away the nebula will again become dark.

Camelopardalis (Cam)

Beta (β) ds. (4.0/8.6, sep. 80.8"). Wide yellow and blue pair (AB). From dark sites the system can be seen to be embedded in a streak of dark nebulosity. A third magnitude 11.8", sep (BC) 14.8", component is visible in moderate apertures.

1 ds. (5.7/6.8, sep. 10.3"). White and pale blue pair.

11 & 12 ds. (5.4/6.5, sep, 108.5") White pair. Easy object for binoculars and small telescopes.

29 ds. 6.5/9.5, sep. 25.1". Yellow primary with fainter pale blue secondary.

Struve (Σ)390 ds. 5.1/9.5, sep. 14.8". White primary with fainter purple secondary.

Struve (Σ)1122 ds. (7.8/7.8, sep. 15.4"). Fine pair of equally bright white stars.

Struve (Σ)1625 ds. 7.3/7.8, sep. 14.4". Almost equal pair of yellowish stars.

Struve (Σ)1694 ds. 5.3/5.8, sep.21.6". Easy pair of whitish stars.

NGC1501 (11.5) pn. Bright, large with a blue tint located about 2 degrees south of the open star cluster NGC1502. Resembles the "Eskimo Nebula" in Gemini. NGC1502 (5.7) oc. A fine open cluster. Extending NW of this cluster is a line of 9th and 10th magnitude stars that form **"Kemble's Cascade"**. A beautiful sight in binoculars or a low power wide field eyepiece on small telescopes.

NGC2403 (8.4) sg. Visible in large binoculars. It lies at a distance of 8m light years and is possibly a member of the M81/M82 group.

IC342 (12.0) sg. Once considered to be a member of the Local Group of galaxies. Its low surface brightness makes it a challenge for moderate apertures. This face on spiral galaxy would be an impressive object if it were further away from the plane of the Milky Way.

Gemini (Gem)

Alpha (α) **Castor** ms. 1.9/2.9 sep 4.0". Close visual pair. However each of these is a spectroscopic binary. A more distant ninth magnitude star (red) forms part of an eclipsing binary system. A fascinating family!

Delta (δ) ds. 3.5/8.2 sep 5.8". Yellow primary with bluish secondary.

Kappa (κ) ds. 3.6/8.1 sep 7.1". Orange-yellow primary with bluish companion.

Lambda (λ) ds. 3.6/10.7 sep 9.6". Blue-white primary with bluish companion.

Σ 1108 (Struve) ds. 6.6/8.3 sep 11..5". Yellow primary with bluish companion.

M35 (5.1) oc. Just visible to the naked eye from dark sites. It is a superb object in telescopes. On its western edge lies the more distant open star cluster IC2158.

NGC2129 (10.2) oc. Located about a degree SW of IC2158.

NGC2266 (9.5) oc. Located about two degrees north of ϵ Gem.

NGC2392 (10.5) pn. "**The Eskimo Nebula**" is a fine planetary nebula located about two degrees SE of δ . The nickname is derived from the appearance of a face surrounded by the hood of a parka.

NGC2420 (8.3) oc. Located about two degrees east of the "Eskimo".

Complete this deep-sky tour of Gemini by locating the open star clusters NGC2355 (9.7) and NGC2395 (7.1).

Lepus (Lep)

Lying beneath Orion Lepus is easily recognized by a quadrilateral of four third magnitude stars and contains a variety of deep-sky objects including one Messier object.

Alpha (α) ds; (2.6,11.1; sep. 35.8").

Beta (β) ds; (3.0/7.5; sep. 2.3").

Gamma (γ) ds. (3.7/6.3, sep. 96.3"). Fine yellow and pale orange pair.

Kappa (κ) ds; (4.5/7.4; sep. 2.6"). White and blue companions.

Iota (ι) ds; (4.5/10.8; sep. 12.7").

NGC1974 sg (11.8). Seen almost edge on.

NGC1904 (**M79**) gc (8.0). A fine globular cluster visible as a fuzzy spot in binoculars. Outer edges begin to resolve in 12" (30cm) telescopes.

IC418 pn (10.7). Very small but bright. Central 10.7 mag star surrounded by pale ring. Use a UHC or OIII filter for best results.

Monoceros (Mon)

This faint and rather indistinct constellation is located between Orion and Canis Minor.

Beta (β) ts. 4.7/5.2/6.1 Sep. AB = 7.3', sep BC = 2.8". Striking triple of bluish white stars.

Epsilon (ϵ) ts. 4.5/6.5 sep. 13.4". Close pair of pale yellow stars. The third mag 12.7 bluish white member is visible in 12"+ apertures.

NGC2244 oc (4.8). Fine open star cluster surrounded by NGC2237-9 "**The Rosette Nebula**" which is best seen using a UHC filter. Shows well in photographs.

NGC2261 en (10v). "**Hubble's Variable Nebula**". Located about 2° southwest of NGC2264 this a fascinating object and well worth monitoring for changes in shape and brightness due to the enveloped variable star R Monocerotis. The triangular wedge appears is almost comet like. Detailed star chart available for telescope owners.

NGC2264 oc + en (4.0) "**The Christmas Tree Cluster**". A fine open cluster with associated nebula that includes "**The Cone Nebula**".

NGC2323 (**M50**) oc (5.9). Superb open cluster.

There are many other open clusters in this area of the Milky Way - NGC's 2215(8.4), 2286(7.5), 2301(6.0), 2335(7.2), 2343(6.7), 2353(7.1) and 2506(7.6).

Orion (Ori)

This constellation dominates the winter skies and because it is so easily recognized forms one of the "key constellations" for finding other winter groupings.

Orion's two brightest stars provide a marked contrast. **Betelgeuse** is distinctly orange in colour. It is a red giant star entering old age. **Rigel** is a brilliant blue/white star indicating the exuberance of youth. Betelgeuse is slightly variable in brightness, range 0.1 - 0.9 and bears the designation α (alpha) indicating that it was brighter than Rigel, β (beta) (0.1), when stars were given these designations. Rigel is now the brighter of the two so either early magnitude estimates were wrong or Betelgeuse has dimmed slightly.

Beta (β) Rigel ds. 0.1/6.8 sep. 9.5". Brilliant bluish white primary with much fainter bluish secondary.

Eta (η) ds. 3.6/5.0 sep. 1.5". Close pair of white stars.

Delta (δ) ds. 2.0/6.9 sep. 52.6". Blue white primary with pale blue secondary.

Lambda (λ) ds. 3.5/5.6 sep. 4.4". White stars. Part of a quadruple system.

Theta-1 (θ) ms. "The Trapezium". AB: 6.7/7.9 sep. 8.8"; CD: 5.1/6.7 sep. 13.4". Superb object!

Iota (ι) ts. 2.8/7.3 sep. 11.3". White primary with pale blue secondary. The third reddish 11th magnitude component is located 50" away.

Sigma (σ) ms. 4.0/10.3 sep. 11.4. Colourful multiple star. Fainter triple star Σ 761 in same field.

Zeta (ζ) Alnitak ds. 1.9/4.0 sep. 2.3". Bluish white stars. Part of a triple system with "**The Flame Nebula**" (NGC2024) in the field of view.

NGC1909 "**The Witch Head Nebula**" about 2½° west of Rigel is a target for imagers.

NGC1976 (**M42**) en. One of the most famous objects in the sky. Marking Orion's sword the "**Great Orion Nebula**" is visible to the naked eye as a faint misty patch. Even a pair of binoculars or small telescope will begin to reveal detail. Increasing aperture and low power bring increasing rewards for the visual observer. Embedded in the nebula is **Theta (θ^1) Ori**. A group of four young stars, mags 5.4, 6.3, 6.8 and 7.0, aptly called "**The Trapezium**". The whole nebula is a stellar nursery with spectacular images being obtained from large Earth based telescopes and the Hubble Space Telescope. M42 is an ideal target for imaging.

NGC1982 (**M43**) en. A small patch of nebulosity on the northern edge of M42.

NGC2024 (en), nicknamed "**The Flame Nebula**", surrounds ζ Ori Alnitak.

IC434 en is a strip of nebulosity extending southwards from ζ Ori Alnitak. The famous "**Horse's Head Nebula**" (Barnard 33) is a small dark intrusion seen dramatically in photographs. It provides one of the biggest challenges to visual observers requiring very dark transparent skies. Responds well to a H-beta nebula filter.

NGC2068 (**M78**) (8.0) rn is a small patch of nebulosity about two degrees NNE of ζ Ori.

NGC2112 (9.1) oc is an open star cluster about two degrees east of M78.

Other open clusters worth locating are NGC2186 (8.7), NGC2169 (5.9) and NGC2175 (6.8) which superimposes a small patch of nebulosity NGC2174.

Long exposure photographs reveal a long arc of nebulosity curving up the east side Orion called "**Barnard's Loop**". It is extremely difficult to discern visually almost regardless of aperture. Remarkably it has been seen with the naked eye (initially by accident!) from dark sites using O III or UHC filters. The "Loop" is a faint ring of hot gas some 14° by 10° with the western part of the ring being less distinct. The "ring" may be due to radiation pressure from the hot young stars in the region of Orion's belt/sword acting on interstellar material. A less favoured school of thought is that it may be a supernova remnant.

Taurus (Tau)

Lambda (λ) vs. Eclipsing binary, range 3.4 to 4.1, period 3.95d.

Phi (ψ) ds. 5.0/8.4 sep 52.1". Fine contrasting deep yellow primary with blue companion.

Σ 495 (Struve) ds. 6.0/8.8 sep 3.8". Fine pair of yellow stars.

Σ 742 ds. 7.1/7.5 sep 4". Fine pair in the same wide field as the Crab Nebula (M1) to the E. Often overlooked when locating M1.

47 Tauri ds. 4.9/7.4 sep 1.1". Very close pair of yellow stars.

T vs. Irregular variable, range 9.4 to 13.5, period erratic.

RV vs. Irregular variable, range 9.5 to 13, period 79d.

RR vs. Irregular variable, range 9.9 to 13, period "chaotic".

M45 - "The Pleiades" or "Seven Sisters" oc. Probably the most famous star cluster. Test your eyesight from a dark site by counting the number of naked eye stars that are visible. Seven should readily be seen. Keen vision will lead you into double figures. A test for moderate apertures is the nebulosity around some of the other brighter stars of the group, especially Merope. Nebula/UHC filters will help.

The Hyades oc. Another naked eye star cluster although more spread out than the Pleiades. At a distance of "a mere" 150 ly it is the closest star cluster to us. The first magnitude star Aldebaran (0.9) is not a member of the cluster and appears brighter as it is closer to us, 21 parsecs - about halfway to the clusters centre. Aldebaran has a faint companion (13.4) separation 30.4". Spotting the companion is not easy even in moderate telescopes.

NGC1514 (10.0) pn. Not an easy object in small instruments. An OIII filter will help.

NGC1554/1555 is a faint variable reflection nebula illuminated by the very young star T Tauri (see above).

NGC1647 (6.4) oc. Rich cluster.

NGC1746 (6.1) oc. Poorer than 1647

NGC1808/1817 (7.0/7.7) oc. a pleasant "double cluster" when observed at low powers.

NGC1952 (**M1**)(8.5) snr. "**The Crab Nebula**". One of the most studied astronomical objects in recent decades. A "new star" appeared in 1054 and over a few months faded from view. Centuries later the faint oval patch was discovered by Dr John Bevis in 1731 and independently by Charles Messier on 12th September 1758 while observing the comet of that year. Its true nature was not recognized until much later. In 1942 astronomers had speculated that a peculiar star in the nebula might be a neutron star. In 1968 radio astronomers discovered that the central star was a pulsar spinning at 30 times a second i.e. the period of the pulses is 33 milliseconds. The "Crab" is also a strong emitter of X-rays.

Happy and Successful Observing!

P.V.H.